

Azimuthal dependence of pion source radii in Pb+Au collisions at 158 A GeV/c

D. Adamová^a, G. Agakichiev^b, A. Andronic^c, D. Antończyk^d, H. Appelshäuser^d, V. Belaga^b,
J. Bielčíková^{e,f}, P. Braun-Munzinger^c, O. Busch^f, A. Cherlin^g, S. Damjanović^f, T. Dietel^h, L. Dietrich^f,
A. Dreesⁱ, W. Dubitzky^f, S. I. Esumi^f, K. Filimonov^f, K. Fomenko^b, Z. Fraenkel^{g,i}, C. Garabatos^c,
P. Glässel^f, G. Hering^c, J. Holeczek^c, M. Kalisky^h, S. Kniege^d, V. Kushpil^a, A. Maas^c, A. Marín^c,
J. Milošević^f, D. Miśkowiec^c, R. Ortega^f, Y. Panebrattsev^b, O. Petchenova^b, V. Petráček^f, M. Płoskoń^d,
S. Radomski^f, J. Rak^c, I. Ravinovich^g, P. Rehak^j, H. Sako^c, W. Schmitz^f, S. Schuchmann^d,
J. Schukraft^k, S. Sedykh^c, S. Shimansky^b, R. Soualah^f, J. Stachel^f, M. Šumbera^a, H. Tilsner^f,
I. Tserruya^g, G. Tsileadakis^c, J. P. Wessels^h, T. Wienold^f, J. P. Wurm^e, S. Yurevich^c, V. Yurevich^b

^a*NPI ASCR, Řež, Czech Republic*

^b*JINR Dubna, Russia*

^c*GSI Darmstadt, Germany*

^d*Frankfurt University, Germany*

^e*MPI, Heidelberg, Germany*

^f*Heidelberg University, Germany*

^g*Weizmann Institute, Rehovot, Israel*

^h*Münster University, Germany*

ⁱ*SUNY Stony Brook, U.S.A.*

^j*BNL, Upton, U.S.A.*

^k*CERN, Geneva, Switzerland*

(Dated: May 16, 2008)

We present results of a two-pion correlation analysis performed with the Au+Pb collision data collected by the upgraded CERES experiment in the fall of 2000. The analysis was done in bins of the reaction centrality and the pion azimuthal emission angle with respect to the reaction plane. The pion source, deduced from the data, is slightly elongated in the direction perpendicular to the reaction plane, similarly as was observed at the AGS and at RHIC.

PACS numbers: 25.75.Gz, 25.75.Ld

I. INTRODUCTION

Two-particle correlations provide unique access to the spatial extension of the source of particles emitted in the course of heavy ion collisions at relativistic energy (for a recent review see [1]). The relation between the experimental correlation function, defined as the relative momentum distribution of pairs, normalized to the analogous distribution obtained via event mixing, and the size of the fireball is especially simple in case of identical pions where the main source of correlations is the Bose-Einstein statistics. In fact, in the pure boson case the correlation function is

$$C(\mathbf{q}, \mathbf{P}) = 1 + \frac{|\int d^4x S(x, P) \exp(i\mathbf{q} \cdot \mathbf{x})|^2}{|\int d^4x S(x, P)|^2} \quad (1)$$

with the source function $S(x, P)$ describing the single particle density in 8-dimensional position-momentum space at freeze-out. The correlation function C depends on the momentum difference $\mathbf{q} = \mathbf{p}_2 - \mathbf{p}_1$ and the mean pion momentum $\mathbf{k} = (\mathbf{p}_1 + \mathbf{p}_2)/2$. The width of the peak

at $\mathbf{q} = 0$ is inversely proportional to the source radius. A particularly exciting prospect is to look for a source asymmetry possibly reflecting the initial asymmetry of the fireball created in collisions with finite impact parameter. Indeed, significant dependence of the source radii on the azimuthal emission angle with respect to the reaction plane, defined by the beam axis and the impact parameter vector, was observed in Au+Au collisions at 2-6 GeV [2] and at $\sqrt{s}=200$ GeV [3]. In this paper we present results of the first analysis of the azimuthal dependence of pion source radii at SPS energies.

II. EXPERIMENT

The CERES/NA45 experiment at the CERN SPS is described in detail in [4]. The upgrade by a radial Time Projection Chamber (TPC) in 1997-1998, in addition to improving the dilepton mass resolution, enhanced the experiment's capability of studying hadronic observables. The cylindrical symmetry of the experiment was preserved during the upgrade, making the setup ideally suited to address azimuthal anisotropies. About 30 million Pb+Au collision events at 158 A GeV/c were collected in the fall of 2000, most of them with centrality within the top 7% of the geometrical cross section σ_G .

[†]Deceased

Small samples of $\sigma/\sigma_G=20\%$ and minimum bias collisions, as well as a short run at 80 A GeV/c, were recorded in addition.

III. DATA ANALYSIS

The results presented here are based on a correlation analysis of the high-statistics Pb+Au data set from the year 2000 [5]. The results of an azimuthal-angle averaged analysis in [5] are consistent with the previously published CERES data [6] and with a recent analysis by the NA49 Collaboration [7]. The main steps of the azimuthal-angle dependent analysis are described below.

A. Event selection

The collision centrality was determined via the charged particle multiplicity around midrapidity $y_{\text{beam}}/2=2.91$. Two variables, the amplitude of the Multiplicity Counter (MC) (single scintillator covering $2.3 < \eta < 3.5$) and the track multiplicity in the TPC ($2.1 < \eta < 2.8$), were alternatively used as the centrality measure. Knowing the data acquisition dead time factor and the target thickness, and assuming that all beam particles were hitting the target, the event counts were translated to the cross section for collisions with a given multiplicity. The centrality variable used in this paper is defined as the integrated cross section divided by the geometrical cross section $\sigma_G=6.94$ barn.

The fireball created in a collision with finite impact parameter is elongated in the direction perpendicular to the reaction plane. In the course of expansion, with the pressure gradient larger in-plane than out-of-plane, the initial asymmetry should get reduced or even reversed. Experimentally, the source eccentricity at the time of decoupling can be determined from an analysis of two-pion correlations as function of $\Phi^* = \Phi_{\text{pair}} - \Phi_{\text{RP}}$, the pair emission angle with respect to the reaction plane.

The azimuthal angle of the reaction plane Φ_{RP} was estimated by the preferred direction of the particle emission (elliptic flow). For this, in each event the flow vector $\mathbf{Q}_2$ was constructed out of the measured particles, characterized by transverse momenta p_t and azimuthal emission angles ϕ [8, 9]:

$$Q_2^X = \frac{1}{N} \sum_{i=1}^N p_{t,i} \cos(2\phi_i) \quad (2)$$

$$Q_2^Y = \frac{1}{N} \sum_{i=1}^N p_{t,i} \sin(2\phi_i). \quad (3)$$

The Q_2^X and Q_2^Y components were calibrated (shifted and scaled) such that the peak in the (Q_2^X, Q_2^Y) distribution was centered at (0,0), and its widths in the directions of X and Y were equal. The reaction plane angle was

calculated (modulo π) from the calibrated $\mathbf{Q}_2$ via

$$\Phi_{\text{RP}} = \frac{1}{2} \arctan \frac{Q_2^Y}{Q_2^X}. \quad (4)$$

The resolution of the so determined reaction plane angle, estimated via the subevent method [8, 10], was $30\text{--}34^\circ$.

The event mixing, needed to obtain the denominator of the correlation functions, was performed in bins of centrality (2% of σ_G), event plane (15°), and vertex (same target disk). This ensures that the shape of the background is identical to that of the signal in all respects except for the femtoscopic correlations.

B. Track selection

Only TPC tracks with at least 12 (out of maximally 20) hits and a reasonably good χ^2 , falling into the fiducial acceptance $0.125 < \theta < 0.240$, were used in the analysis. A momentum dependent dE/dx cut was applied to reduce the contamination of the pion sample. Pions from K^0 and Λ^0 decays were suppressed by a 2.5σ matching cut between the silicon vertex detectors and the TPC.

C. Pair selection

The two-track resolution cuts applied to the true pairs and to the pairs from event mixing were different for the two possible track pair topologies: in the case of the magnetic field bringing the tracks apart from each other (sailor) the required two-track separation was $\Delta\phi > 38\text{--}45$ mrad, depending on the transverse momentum; for the opposite (cowboy) case, $\Delta\phi > 90\text{--}140$ mrad was used. The polar angle cut was the same for both topologies ($\Delta\theta > 8\text{--}9$ mrad). It should be noted that the required two-track cuts depended somewhat on the quality cuts applied to single tracks: the higher the number of hits required for single tracks, the more pairs were lost because of the finite two-track resolution.

D. Correlation functions

The two-pion analysis was performed in the longitudinally co-moving frame (LCMS) defined by the vanishing z -component of the pair momentum. The momentum difference in this frame, $\mathbf{q} = \mathbf{p}_2 - \mathbf{p}_1$, was decomposed into the “out”, “side”, and “long” components following the Bertsch-Pratt convention, with q_{out} pointing along the pair transverse momentum and q_{long} along the beam [11]. The particles were numbered such that q_{side} was always positive.

The $\pi^-\pi^-$ and $\pi^+\pi^+$ correlation functions were fitted

by

$$\begin{aligned} C_2(\mathbf{q}) &= N \cdot \{(1 - \lambda) + \lambda \cdot F_c(\mathbf{q})(1 + G(\mathbf{q}))\}, \\ G(\mathbf{q}) &= \exp\left(-\sum_{i,j} R_{ij}^2 q_i q_j\right) \end{aligned} \quad (5)$$

with the indices $i, j = \{\text{out, side, long}\}$. The normalization factor N was needed because the number of pairs from event mixing was four times higher than of signal pairs. The correlation strength $\lambda < 1$ reflects the contribution of pions from long-lived resonances, the contamination of the pion sample by other particle species, and the finite $\mathbf{q}$ -resolution. The resulting source radii $\sqrt{R_{ij}^2}$ describe the size of the source emitting pions of a given momentum [1]. The $F_c(q_{inv})$ factor, $q_{inv} = \sqrt{-(p_2^\mu - p_1^\mu)^2}$, accounts for the mutual Coulomb interaction between the pions and was calculated by averaging the nonrelativistic Coulomb wave function squared over a realistic source size. The Coulomb factor was attenuated by λ similarly as the rest of the correlation function peak; the importance of this approach was demonstrated in [6]. The fits were performed by the minimum negative log-likelihood method, assuming that the number of true pairs in a given bin is distributed around the expected mean value (equal to the number of mixed pairs times the fit function) according to the Poisson statistics. The source radii obtained from the fit were corrected for the finite momentum resolution

$$\frac{\Delta p}{p} = 2\% \oplus 1\% \cdot p/(\text{GeV}/c). \quad (6)$$

The correction was determined by Monte Carlo and is rather insignificant for R_{side} and R_{long} ; for R_{out} it gets as large as $\approx +20\%$ for the highest pion momenta.

E. Azimuthal dependence of pion source radii

Correlation functions were generated separately for pion pairs with different azimuthal angles with respect to the reaction plane $\Phi^* = \Phi_{\text{pair}} - \Psi_{\text{RP}}$. For this, the pions were sorted into eight bins covering $(-\pi/2, \pi/2)$. The eight correlation functions were fitted [15] and the resulting squared source radii were parametrized by [12]

$$\begin{aligned} R_{\text{out}}^2 &= R_{\text{out},0}^2 + 2R_{\text{out},2}^2 \cos(2\Phi^*) \\ R_{\text{side}}^2 &= R_{\text{side},0}^2 + 2R_{\text{side},2}^2 \cos(2\Phi^*) \\ R_{\text{long}}^2 &= R_{\text{long},0}^2 + 2R_{\text{long},2}^2 \cos(2\Phi^*) \\ R_{\text{os}}^2 &= 2R_{\text{os},2}^2 \sin(2\Phi^*) \\ R_{\text{ol}}^2 &= R_{\text{ol},0}^2 + 2R_{\text{ol},1}^2 \cos(\Phi^*) \\ R_{\text{sl}}^2 &= 2R_{\text{sl},1}^2 \sin(\Phi^*). \end{aligned} \quad (7)$$

While the $R_{\text{out},0}^2$, $R_{\text{side},0}^2$ and $R_{\text{long},0}^2$ obtained represent the phi-averaged squared pion HBT radii, the second Fourier coefficients $R_{i,2}^2$ describe the eccentricity of the

observed pion source. Since the reaction plane is known modulo π odd Fourier components should vanish. If the pion source were to reflect the initial collision geometry (almond shape out-of-plane) a positive $R_{\text{side},2}^2$ and $R_{\text{os},2}^2$ and a negative $R_{\text{out},2}^2$ should be expected. For symmetry reasons all anisotropies should disappear in the limit of central collisions.

The second Fourier coefficients $R_{i,2}^2$ have been corrected for the reaction plane resolution by dividing them by the mean cosine of twice the difference between the reconstructed and the true reaction plane angles $\langle \cos[2(\Psi_{\text{RP}}^{\text{rec}} - \Psi_{\text{RP}})] \rangle$, estimated via the measured mean cosine of the difference between two subevents:

$$R_{i,2}^2 \rightarrow \frac{R_{i,2}^2}{\sqrt{2\langle \cos[2(\Psi_b - \Psi_a)] \rangle}}, \quad (8)$$

similarly as it is done for flow measurements [10]. The correction factor was between 4.8 and 2.4 (centralities of 0-2.5% and 25-70%, respectively). The appropriateness of the flow correction for two-pion correlation radii has been verified with a Monte Carlo simulation of pion pairs emitted from an elliptical source; the observed reduction of the second Fourier coefficients of squared radii was, within the statistical errors of the simulation (about 10%), in agreement with Eq. (8).

F. Systematic errors

The fact that separate analyses of positive and negative pions give consistent results indicates that the limited particle identification is not causing any bias. The systematic error related to the reaction plane resolution correction, based on the comparison between Eq. (8) and a numerical simulation, is estimated to be not larger than 10%. Autocorrelations, i.e. using the same particles to determine the event plane and the HBT radii, might be another source of systematic errors; unfortunately, an attempt to exclude the two particles from the event plane construction led to a strong distortion of the correlation functions as it was not clear how to do the exclusion consistently for the true and mixed pairs. An independent way to estimate the overall systematic errors is by inspecting (see Fig. 1 in the next section) the coefficients $R_{\text{ol},1}^2$ and $R_{\text{sl},1}^2$ which should be zero because the reaction plane is known only modulo π . Averaged over centralities, they are 0.40 (16) and 0.22 (6) fm^2 , respectively. Attributing part (one sigma) of the deviation to statistical fluctuation we are left with a discrepancy of about 0.3 fm^2 which we thus assume to be the systematic uncertainty of the results presented.

IV. RESULTS AND DISCUSSION

The extracted Fourier coefficients are shown in Table I and Fig. 1 [16]. The anisotropies in the out and

TABLE I: Fourier coefficients from a fit of Eq. (7) to the pion source radii squared. The values are in fm².

centrality	mean cent	$R_{\text{out},0}^2$	$R_{\text{side},0}^2$	$R_{\text{long},0}^2$	$R_{\text{ol},0}^2$	$R_{\text{out},2}^2$	$R_{\text{side},2}^2$	$R_{\text{long},2}^2$	$R_{\text{os},2}^2$
0-2.5 %	1.3 %	29.34(10)	24.33(08)	33.10(11)	-6.56(21)	-0.25(25)	-0.03(21)	-1.43(26)	0.19(17)
2.5-5 %	3.7 %	28.01(08)	23.29(06)	31.91(09)	-5.72(18)	-0.56(19)	-0.45(15)	-1.23(19)	0.45(13)
5-7.5 %	6.1 %	26.62(08)	22.04(06)	30.61(10)	-5.22(15)	-0.96(19)	-0.06(15)	-0.59(19)	0.46(12)
7.5-10 %	8.1 %	25.39(17)	21.00(13)	29.22(21)	-5.20(31)	-1.15(32)	-0.11(26)	-1.25(32)	0.63(21)
10-15 %	11.6 %	23.83(19)	19.87(14)	27.31(23)	-4.33(37)	-1.14(35)	0.42(27)	-1.40(34)	0.39(23)
15-25 %	17.5 %	21.35(23)	17.62(17)	24.64(28)	-3.67(48)	-1.03(31)	0.38(24)	-1.27(30)	0.35(21)
25-70 %	30.0 %	14.28(49)	12.29(35)	17.42(61)	-4.23(116)	-0.98(30)	0.36(23)	-1.20(29)	0.34(20)

side directions indicate a pion source elongated out-of-plane. For comparison, the AGS [2] and RHIC [3, 13] values, obtained by performing the fits using Eq. (7) on their published radii (the AGS results were subsequently corrected for their reaction plane resolution), are represented in Fig. 1 by open symbols and stars. It appears that the anisotropies in the out and side directions at SPS energy are rather similar to those observed at RHIC. The geometrical pion source eccentricity can be quantified via $\varepsilon \simeq 2R_{\text{side},2}^2/R_{\text{side},0}^2$ and has, for a




FIG. 1: Azimuthal pion source eccentricity, represented by the second Fourier coefficient of squared radii $R_i^2(\Phi^*)$, measured in Pb+Au collisions as a function of centrality. Positive and negative pion pairs have been combined. The mean pion transverse momentum is 0.23 GeV/c. For comparison, analogous measurements at the AGS (open green symbols) and RHIC (blue diamonds and stars for 130 and 200 GeV, respectively) are shown. The last two panels show the first Fourier coefficients $R_{\text{ol},1}^2$ and $R_{\text{sl},1}^2$ which should be zero by construction. The estimated systematic error is 0.3 fm².

centrality of 15-20%, a value of 0.043(27), significantly less than the initial fireball eccentricity $\varepsilon_{\text{initial}} \approx 0.20$ [3]. The fact that the magnitude of the R_{side} anisotropy seems to be weaker than that of R_{out} indicates that not only the source geometry but also azimuthal dependence of the emission time might play a role. On the other hand, $R_{\text{out},2}^2 - R_{\text{side},2}^2 + 2R_{\text{os},2}^2$, averaged over centralities, is 0.06(18) fm², consistent with zero. This implies that the sum rule from [12], which is supposed to be valid for systems with emission time independent on the azimuthal angle, works rather well.

The R_{long} anisotropy is negative and roughly independent of centrality. Series of checks were performed to make sure that this is not an artefact of the analysis. This result might indicate that R_{long} is sensitive to fluctuations of the azimuthal particle density. Hydrodynamic calculation of central Pb+Au collisions at this energy yields a similar amount of anisotropy in $R_{\text{long},2}^2$ which, however, is centrality dependent [14]. This, and the fact that the hydrodynamic calculation overpredicts the overall magnitude of R_{long} , indicates that the knowledge about the mechanism leading to oscillations of R_{long} is still incomplete.

The source radius anisotropies for centrality 15-20% are shown as a function of the collision energy in Fig. 2. The SPS result fits rather well into the beam energy systematics. The apparent (albeit not statistically significant) fast change of $R_{\text{side},2}^2$ between AGS and SPS and, even more, the negative $R_{\text{long},2}^2$ developing when going down in energy from RHIC to SPS, make the perspective of a systematic study in the course of the low-energy scan at RHIC especially attractive. With the statistical errors of the present AGS data even significant structures in the energy dependence of the pion source anisotropy cannot be excluded.

V. SUMMARY

We have analyzed the azimuthal angle dependence of the pion HBT radii in Pb+Au collisions at the top SPS energy. The source anisotropy in the out and side directions has the same sign and similar magnitude as the one measured at the AGS and at RHIC, and in-




FIG. 2: Collision energy dependence of the pion source anisotropy in Au+Au and Pb+Au collisions at the 15-20% centrality. The meaning of the symbols is the same as in Fig. 1. The SPS results are rather similar to those obtained at RHIC, except for R_{long} which is significantly off-zero. The estimated systematic error is 0.3 fm^2 .

indicates a pion source elongated out-of-plane. The side anisotropy is somewhat smaller than the other which suggests that finite emission times may play a role. The source anisotropy in the long-direction is negative for all centralities indicating that R_{long} might be sensitive to particle density fluctuations.

The CERES collaboration acknowledges the good performance of the CERN PS and SPS accelerators as well as the support from the EST division. We would like to thank R. Campagnolo, L. Musa, A. Przybyla, W. Seipp and B. Windelband for their contribution during construction and commissioning of the TPC and during data taking. We are grateful for excellent support by the CERN IT division for the central data recording and data processing. This work was supported by GSI, Darmstadt, the German BMBF, the German VH-VI 146, the US DoE, the Israeli Science Foundation, the Check Science Foundation contract No. 202/03/0879 and the MINERVA Foundation.

-
- [1] M.A. Lisa, S. Pratt, R. Soltz, U. Wiedemann, Ann.Rev.Nucl.Part.Sci.55(2005)357
 - [2] M.A. Lisa et al, Phys. Lett. B496, 1 (2000).
 - [3] J. Adams et al, Phys. Rev. Lett. 93, 012301 (2004).
 - [4] D. Adamová et al., arXiv:0802.1443v1[nucl-ex], accepted for publication in Nucl. Instr. Meth.
 - [5] D. Antończyk, Ph.D. thesis, Technical University Darmstadt 2006.
 - [6] D. Adamová et al., Nucl. Phys. A714, 124 (2003).
 - [7] S. Kniege, Ph. D. thesis, Frankfurt University 2005; C. Alt et al, arXiv:0709.4507v2 [nucl-ex].
 - [8] P. Danielewicz and G. Odyniec, Phys. Lett. 157B, 146 (1985).
 - [9] J. Barrette et al, Phys. Rev. Lett. 73, 2532 (1994); S. Voloshin and Y. Zhang, Z.Phys.C70, 665 (1996).
 - [10] A.M. Poskanzer and S.A. Voloshin, Phys. Rev. C58, 1671 (1998).
 - [11] G.F. Bertsch, Nucl. Phys. A498, 173c (1989); S. Pratt, Phys. Rev. D33, 1314 (1986).
 - [12] U. Heinz, A. Hummel, M.A. Lisa, U.A. Wiedemann, Phys. Rev. C66, 044903 (2002).
 - [13] R.C. Wells, Ph.D. thesis, Ohio State University 2002.
 - [14] Calculation by P. Huovinen. The paper is in preparation.
 - [15] To avoid problems caused by the non-gaussian shape of inclusive correlation functions the fit was actually performed in bins of pair p_t ; the $R_{i,2}$'s discussed in this paper are weighted averages of the $R_{i,2}$'s obtained for different p_t 's.
 - [16] We refrain in Fig. 1 from normalizing the coefficients to the corresponding mean source radii. In fact, since the second Fourier coefficients measured at RHIC [3] seem to be, within the measurement errors, independent of transverse momentum while the mean radii vary by about 40%, normalization to mean radii could wash out the weak oscillation we are after.